



Regulated Cycling of Mitochondrial Hsp70 at the Protein Import Channel

Qinglian Liu, *et al.*
Science **300**, 139 (2003);
DOI: 10.1126/science.1083379

The following resources related to this article are available online at www.sciencemag.org (this information is current as of May 19, 2009):

Updated information and services, including high-resolution figures, can be found in the online version of this article at:

<http://www.sciencemag.org/cgi/content/full/300/5616/139>

Supporting Online Material can be found at:

<http://www.sciencemag.org/cgi/content/full/300/5616/139/DC1>

This article **cites 16 articles**, 3 of which can be accessed for free:

<http://www.sciencemag.org/cgi/content/full/300/5616/139#otherarticles>

This article has been **cited by** 42 article(s) on the ISI Web of Science.

This article has been **cited by** 19 articles hosted by HighWire Press; see:

<http://www.sciencemag.org/cgi/content/full/300/5616/139#otherarticles>

This article appears in the following **subject collections**:

Cell Biology

http://www.sciencemag.org/cgi/collection/cell_biol

Information about obtaining **reprints** of this article or about obtaining **permission to reproduce this article** in whole or in part can be found at:

<http://www.sciencemag.org/about/permissions.dtl>

Distinct categories of death stimuli require BAX and BAK at either mitochondria or ER control points (fig. S7). BAX and BAK at mitochondria are requisite for "BH3-only" proteins, whereas regulation of ER Ca^{2+} by BAX and BAK proved an obligate control point for lipid second messengers (17–19) and oxidative stimuli (31). A third category of stimuli is influenced by control points at both mitochondria and ER and includes intrinsic signals initiated by staurosporine, etoposide, and brefeldin-A. The coordinate roles for BAX and BAK at the ER and mitochondria illustrate the importance of Ca^{2+} dynamics between these organelles in apoptosis.

References and Notes

- D. R. Green, J. C. Reed, *Science* **281**, 1309 (1998).
- K. F. Ferri, G. Kroemer, *Nature Cell Biol.* **3**, E255 (2001).
- J. M. Adams, S. Cory, *Trends Biochem. Sci.* **26**, 61 (2001).
- M. C. Wei et al., *Science* **292**, 727 (2001).
- M. C. Wei et al., *Genes Dev.* **14**, 2060 (2000).
- E. H. Cheng et al., *Mol. Cell* **8**, 705 (2001).
- P. Li et al., *Cell* **91**, 479 (1997).
- R. Rizzuto et al., *Science* **280**, 1763 (1998).
- G. Hajnoczky, L. D. Robb-Gaspers, M. B. Seitz, A. P. Thomas, *Cell* **82**, 415 (1995).
- R. Rizzuto, A. W. Simpson, M. Brini, T. Pozzan, *Nature* **358**, 325 (1992).
- P. Nicotera, S. Orrenius, *Cell Calcium* **23**, 173 (1998).
- P. Bernardi, *Physiol. Rev.* **79**, 1127 (1999).
- D. Decaudin et al., *Cancer Res.* **57**, 62 (1997).
- M. Lam et al., *Proc. Natl. Acad. Sci. U.S.A.* **91**, 6569 (1994).
- P. Pinton et al., *J. Cell Biol.* **148**, 857 (2000).
- R. Foyouzi-Youssefi et al., *Proc. Natl. Acad. Sci. U.S.A.* **97**, 5723 (2000).
- L. M. Obeid, C. M. Linardic, L. A. Karolak, Y. A. Hannun, *Science* **259**, 1769 (1993).
- G. Szalai, R. Krishnamurthy, G. Hajnoczky, *EMBO J.* **18**, 6349 (1999).
- L. Scorrano, D. Penzo, V. Petronilli, F. Pagano, P. Bernardi, *J. Biol. Chem.* **276**, 12035 (2001).
- D. M. Hockenbery, Z. N. Oltvai, X. M. Yin, C. L. Millman, S. J. Korsmeyer, *Cell* **75**, 241 (1993).
- B. Gajkowska, T. Motyl, H. Olszewska-Badarczuk, M. M. Godlewski, *Cell Biol. Int.* **25**, 725 (2001).
- L. K. Nutt et al., *J. Biol. Chem.* **277**, 20301 (2002).
- L. S. Jouaville, F. Ichas, E. L. Holmuhamedov, P. Camacho, J. D. Lechleiter, *Nature* **377**, 438 (1995).
- E. Boitier, R. Rea, M. R. Duchon, *J. Cell Biol.* **145**, 795 (1999).
- A. Weiss, J. B. Imboden, *Adv. Immunol.* **41**, 1 (1987).
- S. A. Oakes, J. T. Opferman, unpublished data.
- F. Vanden Abeele et al., *Cancer Cell* **1**, 169 (2002).
- C. Li et al., *Proc. Natl. Acad. Sci. U.S.A.* **99**, 9830 (2002).
- K. K. Wang, *Trends Neurosci.* **23**, 20 (2000).
- T. Nakagawa, J. Yuan, *J. Cell Biol.* **150**, 887 (2000).
- T. Finkel, N. J. Holbrook, *Nature* **408**, 239 (2000).
- We thank A. Vukovic and C. Ong for technical assistance, and E. Smith for editorial assistance. L.S. is a Human Frontier Science Program fellow and a Dulbecco-Telethon Institute Assistant Scientist. Supported by NIH training grant T32HL07627 (S.A.O.), a postdoctoral fellowship (DRG 1664) from the Damon Runyon Cancer Research Foundation (J.T.O.), and NIH grant R37CA50239.

Supporting Online Material

www.sciencemag.org/cgi/content/full/1081208/DC1
Materials and Methods
Figs. S1 to S7
Movies S1 to S4

5 December 2002; accepted 7 February 2003
Published online 6 March 2003;
10.1126/science.1081208
Include this information when citing this paper.

Regulated Cycling of Mitochondrial Hsp70 at the Protein Import Channel

Qinglian Liu,^{1,2*} Patrick D'Silva,^{1*} William Walter,¹ Jaroslaw Marszalek,^{1†} Elizabeth A. Craig^{1§}

Hsp70 of the mitochondrial matrix (mtHsp70) provides a critical driving force for the import of proteins into mitochondria. Tim44, a peripheral inner-membrane protein, tethers it to the import channel. Here, regulated interactions were found to maximize occupancy of the active, adenosine 5'-triphosphate (ATP)-bound mtHsp70 at the channel through its intrinsic high affinity for Tim44, as well as through release of adenosine diphosphate (ADP)-bound mtHsp70 from Tim44 by the cofactor Mge1. A model peptide substrate rapidly released mtHsp70 from Tim44, even in the absence of ATP hydrolysis. In vivo, the analogous interaction of translocating polypeptide would release mtHsp70 from the channel. Consistent with the ratchet model of translocation, subsequent hydrolysis of ATP would trap the polypeptide, driving import by preventing its movement back toward the cytosol.

Because only a few of the hundreds of proteins in mitochondria are encoded by mitochondrial DNA, mitochondrial function depends on efficient import of proteins translated on cytosolic ribosomes. Translocation of these proteins across the inner mitochondrial membrane

through the inner membrane (TIM) channel is driven by an import motor. A critical component of this machine is the major Hsp70 molecular chaperone of mitochondria (mtHsp70; Ssc1 in yeast) (1, 2), which binds short hydrophobic segments of the incoming polypeptide chains. MtHsp70 is tethered to the TIM channel by interaction with the peripherally associated membrane protein Tim44 (3–5). How this interaction promotes the function of mtHsp70 in mitochondrial import is unresolved and remains controversial (6–8). Because the mechanism of mitochondrial import has been addressed primarily by using complex in organellar systems, we purified mtHsp70 and Tim44, thus allowing a dissection of interactions in vitro.

To assess the mtHsp70-Tim44 interaction in vitro, we monitored complex formation by the

ability of Tim44 to be precipitated by mtHsp70-specific antibodies (9). mtHsp70, like other Hsp70s, is an adenosine triphosphatase (ATPase), whose chaperone activity is dependent upon alternating between ATP- and ADP-bound states (10, 11). To mimic the in vivo situation, we carried out reactions in the presence of ATP or ADP. About 50% of the Tim44 was coimmunoprecipitated in the presence of either nucleotide (Fig. 1A; fig. S2). This lack of nucleotide dependence of complex formation was surprising, because coimmunoprecipitation analysis with mitochondrial lysates had indicated that the mtHsp70-Tim44 interaction was extremely ATP sensitive (12–14) (Fig. 1A and fig. S2). To ensure that the interaction of purified Tim44 and mtHsp70 observed in the presence of ATP was not due to hydrolysis during the course of the experiment, we tested the interaction of mtHsp70 with a peptide substrate, an interaction known to be sensitive to ATP both in vivo and in vitro. A model peptide tagged with fluorescein was used to monitor binding in the mtHsp70 peptide-binding cleft under the conditions used to analyze the mtHsp70-Tim44 interaction. As expected from previous results (11), mtHsp70 had a dissociation constant (K_d) of 0.3 μM for peptide in the presence of ADP, but a drastically reduced affinity (~50-fold less) in the presence of ATP or a nonhydrolyzable ATP analog, AMP-PNP (5'-adenylylimido diphosphate) (Fig. 1B). The reduced peptide affinity in the presence of ATP or AMP-PNP indicated that ATP hydrolysis was inconsequential under these assay conditions.

To assess more quantitatively the mtHsp70-Tim44 interaction, we incubated increasing concentrations of mtHsp70 (0.35 to 2.8 μM) in the presence of 0.09 μM Tim44 with ADP, ATP, or AMP-PNP (Fig. 1, C and D). Concentration-dependent binding was observed in all cases,

¹Department of Biochemistry, ²Graduate Program in Biomolecular Chemistry, University of Wisconsin-Madison, Madison, WI 53706, USA.

*These authors contributed equally to this work.

†Present address: Department of Biochemistry and Biophysics, Columbia University, 650 West 168th Street, New York, NY 10032, USA.

‡Present address: Department of Molecular and Cellular Biology, Faculty of Biotechnology, University of Gdansk, 80-822, Gdansk, Klakki 24, Poland.

§To whom correspondence should be addressed. E-mail: ecraig@wisc.edu

REPORTS

with 50% of Tim44 binding occurring at mtHsp70 concentrations between 0.5 and 1.5 μM . Thus, mtHsp70 and Tim44 formed a complex in the presence of ATP, ADP, or AMP-PNP. Because the amount of mtHsp70 required to form a complex was similar in the presence of ATP and AMP-PNP, hydrolysis of ATP was not required for this interaction. In addition, the mtHsp70-Tim44 interaction was fundamentally distinct from a peptide substrate interaction, because mtHsp70 had a similar affinity for Tim44 regardless of its nucleotide state.

In vivo, the critical functional interaction is the formation of a ternary complex between Tim44-bound mtHsp70 and the translocating polypeptide chain at the import channel. MtHsp70(ATP) initiates this interaction because the on rate is very rapid (10). Subsequent hydrolysis of ATP to ADP, which is stimulated by polypeptide binding, stabilizes this interaction, because the off rate of polypeptide is slow in the ADP-bound state. To mimic this interaction in vitro, we added peptide to a preformed mtHsp70-Tim44 complex. The addition of peptide destabilized the mtHsp70(ATP)-Tim44 complex, but not the mtHsp70(ADP)-Tim44 complex (Fig. 2A). Thus, in vivo, the binding of polypeptide to Hsp70 may trigger the activation of the import machine, stimulating the release of mtHsp70 from Tim44.

The results of this in vitro analysis also provide an explanation for the sensitivity of the mtHsp70-Tim44 complex to ATP, but not to ADP, observed in mitochondrial lysates. Lysates contain a variety of polypeptides that may have exposed hydrophobic amino acids, providing mtHsp70 binding sites. The availability of such

substrates would promote dissociation of the mtHsp70(ATP)-Tim44 complex, which otherwise would remain stable. On the other hand, the presence of such potential substrates would be expected to have no effect on the mtHsp70(ADP)-Tim44 complex.



Because polypeptide binding stimulated ATP hydrolysis, we examined whether peptide-induced release of mtHsp70 from Tim44 required hydrolysis by substituting AMP-PNP for ATP. Release still occurred (Fig. 2A). Because the order of Ssc1's release from Tim44 and ATP hydrolysis has mechanistic implications, we compared the rates of the two events (9). Fifteen seconds after addition of peptide, no detectable Tim44-mtHsp70 complex remained (Fig. 2B). The rate of hydrolysis of ATP in a preformed Tim44-mtHsp70(ATP) complex was stimulated by the addition of peptide. However, 15 s after addition of peptide, less than 5% of the ATP was hydrolyzed (Fig. 2B). Thus, ATP hydrolysis was not required for, nor did it occur prior to, release.

Although events subsequent to mtHsp70 binding to translocating polypeptide are critical, efficient import will only occur if occupancy of the productive, ATP-bound form of mtHsp70 at the import channel is maximized. Given that the estimated concentrations of mtHsp70 and Tim44 in mitochondria are in the range of 70 and 3 μM , respectively (15, 16), most of the Tim44 should be complexed with mtHsp70 in vivo. However, barring the effects of unknown regulatory mechanisms, mtHsp70(ATP) and mtHsp70(ADP) would bind with similar affinity. In addition, because mtHsp70s have an intrinsic ATPase activity, conversion of mtHsp70(ATP)-Tim44 to mtHsp70(ADP)-Tim44 could occur at the

channel. Thus, we investigated whether other factors functioned to maximize occupancy by mtHsp70(ATP).

In organello studies indicated that Mge1, a mtHsp70 nucleotide release factor, promotes release of mtHsp70 from Tim44 (17). We examined the effect of Mge1 on the mtHsp70-Tim44 interaction in vitro. Consistent with previous results, Mge1 stimulated the release of mtHsp70(ADP) from Tim44 (Fig. 2; fig. S3). However, Mge1 had little effect on the mtHsp70(ATP)-Tim44 interaction. Thus, Mge1 may facilitate the dissociation of nonproductive mtHsp70(ADP)-Tim44 complexes without affecting the stability of productive mtHsp70(ATP)-Tim44 complexes. In vivo, Mge1 would function to maintain mtHsp70(ATP) at the import channel poised to bind an incoming polypeptide chain.

Next we tested whether a mtHsp70-polypeptide complex recently released from the import channel would compete with



mtHsp70(ATP) for binding to Tim44. Purified Tim44 was added to preformed mtHsp70(ADP)-peptide complex (Fig. 2A). No Hsp70(ADP)-Tim44 complex was detected. Likewise, preformed mtHsp70(ADP)-Tim44 complex was stable in the presence of peptide (Fig. 2A). To determine if a stable triple complex of mtHsp70, Tim44, and peptide could be detected, we incubated the two proteins and fluorescently labeled peptide in various combinations in the presence of ADP, and then separated the complexes by size exclusion chromatography. Whereas mtHsp70-peptide and mtHsp70-Tim44 complexes were detected by this method, no mtHsp70-Tim44-peptide complex was observed (Fig. 3A), suggesting that mtHsp70(ADP) did not bind Tim44 and polypeptide simultaneously.


The results reported here can be incorporated into a comprehensive model of mitochondrial protein import. The functional ATP-bound form of mtHsp70, which can rapidly bind to the incoming polypeptide chain, predominates at the import channel because of its inherent high af-

finity for Tim44, as well as the dissociation of any ADP-bound mtHsp70 by Mge1 (Fig. 3B, point 1). As the presequence of the polypeptide emerges into the matrix driven by the membrane potential, it interacts with mtHsp70. This binding of polypeptide stimulates the rapid release of mtHsp70 from Tim44 (point 2). Polypeptide binding also stimulates ATP hydrolysis (point 3), which stabilizes mtHsp70's interaction with the polypeptide, preventing retrograde movement of the polypeptide back toward the cytosol. The release of mtHsp70 not only allows diffusion of the polypeptide chain, with mtHsp70 attached, away from the import channel, it also frees Tim44 for binding to another mtHsp70, initiating another cycle of mtHsp70 interaction with the polypeptide and driving import of the chain. The recently released mtHsp70-polypeptide complex is not able to rebound to Tim44, and thus it does not compete with binding of free ATP-bound mtHsp70 (point 4). In time, Mge1, acting as a nucleotide release factor (18, 19), binds mtHsp70(ADP), stimulating release of ADP, and thus carrying out its second function in mitochondrial import.

Subsequent binding of ATP causes release of mtHsp70 from the chain, owing to the rapid off rate in this nucleotide state, completing the cycle of mtHsp70-polypeptide interaction. The active, ATP-bound form of mtHsp70 is regenerated, and folding of the polypeptide chain proceeds.

A particularly controversial issue has been whether mtHsp70 functions as a ratchet, allowing movement of the polypeptide chain into the matrix but not back toward the cytosol (Brownian ratchet model), or whether mtHsp70 exerts a force on the chain "pulling" it into the matrix (power-stroke/pulling model) (6-8, 20, 21). According to the power-stroke model, mtHsp70 exerts a force on the polypeptide chain, using Tim44 as a fulcrum as it undergoes a conformational change associated with ATP hydrolysis, which also stabilizes the mtHsp70-polypeptide interaction. The power-stroke model is not compatible with the results presented here without evoking unknown components. Not only is ATP hydrolysis unnecessary for release of mtHsp70 from Tim44, but Tim44 release also is more rapid than the rate of ATP hydrolysis. Our data are completely consistent with the ratchet model.

Fig. 3. Interaction of mtHsp70 with cofactors and substrate peptide. **(A)** Analysis of mtHsp70 complexes by size exclusion chromatography. Reactions: (1) 15 μ g of Tim44 (triangles); (2) 30 μ g of mtHsp70 and fluorescein-labeled peptide P5 (circles); (3) 30 μ g of mtHsp70 incubated with 15 μ g of Tim44 for 30 min, followed by the addition of fluorescein-labeled P5 and further incubation for 1 hour (squares). Reactions were subjected to chromatography on a Superdex 200HR-10/30 column. Fractions were subjected to electrophoresis and immunoblot analysis with mtHsp70- and Tim44-specific antibodies; signals were quantified by densitometry. P5 peptide in each fraction was monitored by fluorescence intensity. Tim44 forms a dimer (16), hence it shows an elution profile similar to that of mtHsp70. Note that when fluorescein-tagged peptide was added to mtHsp70 and Tim44, no fluorescence was detected in the fractions containing the faster moving Tim44-mtHsp70 complex. **(B)** Model of regulation of the interactions between mtHsp70 and Tim44, translocating polypeptide, and Mge1. Circled numbers refer to points of discussion in the text.



References and Notes

1. P. J. Kang *et al.*, *Nature* **348**, 137 (1990).
2. P. Scherer, U. Krieg, S. Hwang, D. Vestweber, G. Schatz, *EMBO J.* **9**, 4315 (1990).
3. A. C. Maarse, J. Blom, L. A. Grivell, M. Meijer, *EMBO J.* **11**, 3619 (1992).
4. P. E. Scherer, U. C. Manning-Krieg, P. Jenö, G. Schatz, M. Horst, *Proc. Natl. Acad. Sci. U.S.A.* **89**, 11930 (1992).
5. C. Weiss *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* **96**, 8890 (1999).
6. A. Matouschek, N. Pfanner, W. Voos, *EMBO Rep.* **5**, 404 (2000).
7. N. Pfanner, A. Geissler, *Nat. Rev. Mol. Cell. Biol.* **5**, 339 (2001).
8. W. Neupert, M. Brunner, *Nat. Rev. Mol. Cell. Biol.* **8**, 555 (2002).
9. Materials and Methods are available as supporting material on Science Online.
10. B. Bukau, A. L. Horwich, *Cell* **92**, 351 (1998).
11. Q. Liu, J. Krzewski, K. Liberek, E. A. Craig, *J. Biol. Chem.* **276**, 6112 (2001).
12. H.-C. Schneider *et al.*, *Nature* **371**, 768 (1994).
13. O. von Ahsen, W. Voos, H. Henninger, N. Pfanner, *J. Biol. Chem.* **270**, 29848 (1995).
14. M. Horst, W. Oppliger, B. Feifel, G. Schatz, B. Glick, *Protein Sci.* **5**, 759 (1996).
15. C. Voisine, W. Walter, E. A. Craig, unpublished results.
16. F. Moro, C. Sirrenberg, H.-C. Schneider, W. Neupert, M. Brunner, *EMBO J.* **18**, 3667 (1999).
17. H.-C. Schneider, B. Westermann, W. Neupert, M. Brunner, *EMBO J.* **15**, 5796 (1996).
18. B. Miao, J. E. Davis, E. A. Craig, *J. Mol. Biol.* **265**, 541 (1997).
19. P. J. Dekker, N. Pfanner, *J. Mol. Biol.* **270**, 321 (1997).
20. C. Voisine *et al.*, *Cell* **97**, 565 (1999).
21. K. Okamoto *et al.*, *EMBO J.* **21**, 3659 (2002).
22. This work was supported by NIH grant RO1GM27870. We thank T. Donohue, P. Kiley, and C. Wiese for helpful comments.

Supporting Online Material
www.sciencemag.org/cgi/content/full/300/5616/139/DC1
 Materials and Methods
 Figs. S1 to S3
 References

12 February 2003; accepted 6 March 2003